



Barnet Products Corporation 140 Sylvan Avenue Englewood Cliffs NJ 07632 Tel 201 346 4620 Fax 201 346 4333 Web barnetproducts.com

BV-OSC

- . Whitening now tested in vivo
- Eliminates Age Spots
- Anti-Oxidant
- · Collagen Synthesis & Protection
- DNA Protection





DESCRIPTION

BV-OSC (Tetrahexyldecyl Ascorbate) is a stable, oil soluble form of Vitamin C.

PROPERTIES

- * BV-OSC at 0.1% reduces melanin synthesis by 80%.
- * BV-OSC at 10% eliminates age spots in 16 weeks.
- * BV-OSC at 3% in vivo reduces Delta-L value by 15% vs. placebo (22 people), a way to measure whitening effect
- * BV-OSC at 0.1% in vitro increases collagen by 50%.
- * BV-OSC tested at 10% in vivo to treat acne with 80% of patients (12 people) satisfied with results
- * BV-OSC increases collagen synthesis at least twice as much as ascorbic acid.
- * BV-OSC inhibits MMP-2 and MMP-9 over 3 times better than ascorbic acid.
- * BV-OSC penetrates the skin 4 times better than Magnesium Ascorbyl Phosphate.
- * BV-OSC delivers pure Vitamin C 50 times better than ascorbic acid.
- * BV-OSC decreases 8-OHdG induced by UV-A.
- * BV-OSC decreases p53 expression induced by UV-B.
- * BV-OSC protects the cells against UV-B better than other esters of Vitamin C.
- * BV-OSC works synergistically with Thiotaine as both penetrate the cells for anti-oxidant activity.

FORMULATION

BV-OSC is a slight to pale yellow liquid with a faint characteristic odor. It is very soluble in ethanol, hydrocarbons, esters and vegetable oils. It is insoluble in glycerin and butylene glycol. BV-OSC should be added into the oil phase at temperatures below 80°C. It can be used in formulas with a pH range of 3 to 6. Use level is 0.5% - 3%. BV-OSC is approved as a quasi-drug in Korea at 2%, and in Japan at 3%.

LEGISLATION

INCI Name: Tetrahexyldecyl Ascorbate

JSCI: 532297 (Ascorbyl Tetra 2-Hexyldecanoate)

CAS: 183476-82-6

EINECS:

KOREA: Approved as a functional ingredient for whitening at 2%.

BV-OSC


BV-OSC ELIMINATES AGE SPOTS

BV-OSC at 10% concentration was applied to the skin of 10 people for a period of 16 weeks. Results (below) show that BV-OSC eliminated age spots.



COMPARISON OF ABILITY FOR COLLAGEN SYNTHESIS

Proline involved in collagen synthesis was labeled by ³H and added to human dermal fibroblasts (NHDF) with various concentrations of BV-OSC/L-ascorbic acid and cultivated for 24 hours. Then collagen fractions were obtained. The amount of ³H taken into the collagen fraction was measured by using a liquid scintillation counter and slot blotter. As shown below, BV-OSC significantly promoted collagen synthesis.



The information contained in this technical bulletin is, to the best of our knowledge, true and accurate. No warranty, expressed or implied is made or intended. The use should be based upon the customer's own investigations and appraisal. No recommendation should be construed as an inducement to use a material in infringement of patents or applicable government regulations.